

Impact Of Global Climate Change On Coastal Ecosystems: An International Study

Dr. Manisha Suhas Ramteke
MA (Geog.) B.Ed. NET, Ph.D.

Abstract

Coastal ecosystems are among the most productive, diverse and socially valuable components of the Earth system, yet they are increasingly exposed to interacting pressures arising from global climate change and local human activities. This research paper examines the international dimensions of climate-change impacts on mangrove forests, coral reefs, seagrass meadows, salt marshes, estuaries, sandy beaches and associated coastal communities. Using a qualitative comparative review of authoritative international assessments and recent peer-reviewed literature, the study analyses five major pathways of impact: sea-level rise, ocean warming and marine heatwaves, ocean acidification, changes in storms and precipitation, and shifts in salinity, sediment and freshwater regimes. Evidence indicates that climate change is not acting as a single stressor; instead, it interacts with coastal development, pollution, overfishing, habitat fragmentation and other anthropogenic pressures. The consequences include coral bleaching, wetland loss, coastal erosion, biodiversity redistribution, reduced fisheries productivity, declining natural shoreline protection and growing risks to livelihoods and infrastructure. International examples from the Great Barrier Reef, the Sundarbans, the Mekong Delta, Caribbean coral-reef systems and small island states illustrate how exposure and adaptive capacity vary geographically. The paper further evaluates ecosystem-based adaptation, blue-carbon conservation, restoration, integrated coastal-zone management, climate-resilient fisheries and international cooperation. The analysis concludes that effective coastal resilience requires simultaneous mitigation of greenhouse-gas emissions and locally appropriate adaptation that protects ecological processes, restores natural buffers and includes coastal communities in decision-making. The paper highlights the importance of transboundary governance, long-term monitoring, equitable finance and science-based planning for sustaining coastal ecosystem services under a changing climate.

Keywords: climate change; coastal ecosystems; sea-level rise; coral reefs; mangroves; seagrass; salt marshes; ocean acidification; marine heatwaves; adaptation; blue carbon; coastal resilience.

1. Introduction

Coastal ecosystems occupy the interface between land and sea and include a diverse mosaic of mangrove forests, coral reefs, seagrass meadows, tidal marshes, estuaries, lagoons, dunes, beaches and shallow marine habitats. These ecosystems support biodiversity and provide services that are fundamental to human societies, including fisheries production, shoreline stabilization, carbon storage, nutrient cycling, tourism, recreation and cultural values. Their significance is global because coastal populations, cities, infrastructure and economic activities are highly concentrated near the sea. The United Nations notes that hundreds of millions of people live in low-lying coastal areas and that climate-driven changes in the ocean increasingly affect food security, livelihoods and biodiversity (United Nations, n.d.).

Climate change has altered the physical and chemical conditions on which coastal ecosystems depend. Global mean sea level has risen substantially since the beginning of the twentieth century and the rate of rise has accelerated. Ocean warming has increased the frequency and intensity of marine heatwaves, while increasing atmospheric carbon dioxide has changed seawater chemistry and contributed to ocean acidification. These changes occur simultaneously with tropical cyclones, heavy rainfall, drought, coastal subsidence and altered river discharge. The IPCC concludes that coastal ecosystems are already affected by the combination of sea-level rise, climate-related ocean changes and non-climatic human pressures (IPCC, 2019; IPCC, 2022).

The scientific challenge is therefore not simply to identify whether climate change affects coasts, but to understand how multiple drivers interact, why vulnerability differs among regions, and which adaptation measures can preserve ecological functions while supporting human well-being. This paper addresses that challenge through an international comparative perspective. It focuses on the ecological mechanisms of change, social and economic consequences, selected regional cases and the policy options available at local, national and international scales.

2. Objectives of the Study

To examine the principal mechanisms through which global climate change affects coastal ecosystems.

To assess ecological, economic and social consequences for coastal communities and ecosystem services.

To compare climate-related coastal risks across selected regions and ecosystem types.

To evaluate adaptation, conservation and restoration approaches, including nature-based solutions and blue-carbon strategies.

To identify major gaps in governance, finance, monitoring and research relevant to international coastal resilience.

3. Research Questions

- What are the major pathways through which global climate change is altering coastal ecosystems?

How do sea-level rise, ocean warming, acidification and extreme events interact with local anthropogenic pressures?

Why do the impacts and adaptive capacities of coastal ecosystems differ among regions?

Which ecosystem-based and policy responses can reduce ecological and social vulnerability?

What forms of international cooperation are required to strengthen long-term coastal resilience?

4. Methodology

The study uses a qualitative, comparative research design based primarily on secondary data. The evidence base combines major assessments by the Intergovernmental Panel on Climate Change (IPCC), United Nations agencies and recent peer-reviewed literature on coastal wetlands, mangroves, coral reefs and climate adaptation. Priority was given to sources that synthesize evidence across regions or provide internationally recognized assessments. The analysis is organized around a driver–impact–response framework: climate driver, ecosystem response, consequences for ecosystem services and communities, and potential adaptation or mitigation response.

Five illustrative regional contexts are considered: the Great Barrier Reef in Australia; the Sundarbans in Bangladesh and India; the Mekong Delta in Viet Nam; Caribbean coral-reef systems; and small island developing states in the Pacific and Indian Ocean. These cases were selected to represent different ecosystem types, climate exposures and governance settings rather than to provide a statistically representative global sample. The study is therefore interpretive and comparative rather than a meta-analysis. A limitation is that coastal responses vary strongly at local scales because of geomorphology, sediment supply, land subsidence, land use, management quality and socioeconomic conditions.

5. Conceptual Framework: Climate Change and Coastal Ecosystem Vulnerability

Coastal vulnerability can be understood through the interaction of exposure, sensitivity and adaptive capacity. Exposure refers to the degree to which an ecosystem or community encounters hazards such as sea-level rise, heatwaves or storms. Sensitivity describes the extent to which ecological structure and functions respond to those hazards. Adaptive capacity concerns the ability of ecosystems and societies to absorb disturbance, reorganize, migrate, restore functions or alter management practices. This framework is especially useful because the same climate driver can produce different outcomes in different locations.

For example, mangroves can tolerate periodic flooding and can sometimes accumulate sediment vertically as sea level rises. However, their adaptation potential is constrained where roads, embankments, cities or other infrastructure prevent landward migration. The IPCC identifies this 'coastal squeeze' as a major factor reducing the capacity of coastal wetlands to respond to rising seas. Similarly, coral reefs can provide wave attenuation and habitat, but warming, acidification, pollution and physical damage reduce coral growth and recovery, weakening their capacity to keep pace with sea-level rise (IPCC, 2019; IPCC, 2022).

The framework also emphasizes cumulative and compound impacts. A marine heatwave can cause coral bleaching; if bleaching is followed by a cyclone, sediment runoff or disease, recovery can be delayed. In estuaries, warming can interact with nutrient pollution to intensify hypoxia. Sea-level rise can combine with storm surge and land subsidence to create much greater flood exposure than any single factor alone. Consequently, coastal climate policy should address interacting risks rather than treating each hazard separately.

6. Major Impacts of Global Climate Change on Coastal Ecosystems

6.1 Sea-Level Rise and Coastal Inundation

Sea-level rise is one of the most consequential long-term climate risks for coastal ecosystems. It increases the frequency of high-water events, coastal flooding and erosion, while altering salinity and the position of tidal zones. The IPCC reports that global mean sea level has risen by approximately 0.20 metres since 1901 and that the rate has accelerated (IPCC, 2022). The consequences are especially significant for low-lying islands, deltas and coastal wetlands.

Coastal wetlands may respond by building elevation through sediment accumulation and organic matter production, but this capacity depends on local sediment supply, tidal range, vegetation productivity and the rate of sea-level rise. Where coastal development blocks inland migration, wetlands can become trapped between rising water and fixed infrastructure. The resulting habitat loss reduces nursery grounds, carbon storage and natural coastal protection. The IPCC assesses that

many coastal ecosystems will face increasing difficulty keeping pace with sea-level rise, with risks escalating under higher-emissions pathways (IPCC, 2022).

6.2 Ocean Warming and Marine Heatwaves

Ocean warming affects metabolism, reproduction, species distributions and ecological interactions. Marine heatwaves are particularly damaging because they can expose organisms to temperatures beyond their physiological tolerance for days or weeks. IPCC assessments identify marine heatwaves as a driver of abrupt changes in community composition, biodiversity loss and disruptions to fisheries and aquaculture (IPCC, 2022).

Coral reefs provide the clearest example. When thermal stress causes corals to expel their symbiotic algae, bleaching occurs. Severe or repeated bleaching can result in mortality and reduced reef-building capacity. The IPCC estimates that warm-water coral reefs are projected to decline by 70–90% at 1.5°C of global warming and by more than 99% at 2°C, illustrating the narrow thermal margin within which reef ecosystems can persist (IPCC, 2018). The ecological consequences extend to tourism, fisheries, shoreline protection and cultural values.

6.3 Ocean Acidification

The ocean absorbs a substantial fraction of anthropogenic carbon dioxide. This uptake changes seawater carbonate chemistry and lowers pH, reducing the availability of carbonate ions needed by many calcifying organisms. Coral reefs, shellfish and other calcifying organisms can therefore face reduced calcification and greater vulnerability to erosion. Acidification does not act independently; warming, deoxygenation, storms and pollution can compound biological stress. The IPCC identifies coral reefs and other calcified coastal organisms as particularly sensitive to these interacting pressures (IPCC, 2019; IPCC, 2022).

6.4 Changes in Storms, Rainfall and Extreme Events

Tropical cyclones, storm surges, extreme rainfall and flooding can physically restructure coastal habitats. Mangroves and salt marshes may be damaged by severe storms, while beaches and dunes can experience rapid erosion. Coral reefs can suffer breakage and sedimentation. At the same time, extreme rainfall can increase freshwater discharge and nutrient and sediment loading into estuaries and reefs. Climate change therefore alters not only mean environmental conditions but also the frequency, intensity and sequencing of disturbances.

Extreme events can also interact with sea-level rise. A higher baseline sea level means that a storm surge begins from a higher water level, increasing the probability that a previously rare coastal flood becomes a recurrent event. IPCC assessments indicate that historically rare extreme sea-level events are projected to become much more frequent at many locations, increasing risks to ecosystems, settlements and infrastructure (IPCC, 2019; IPCC, 2022).

6.5 Salinity, Freshwater and Sediment Changes

Deltas and estuaries depend on a balance among freshwater inflow, sediment delivery, tides and marine processes. Climate change can modify precipitation, drought frequency, river discharge and sea level, while dams and water extraction can independently reduce sediment and freshwater delivery. Salinization can shift plant communities and affect agricultural land, freshwater supplies and estuarine species. The combined effects are particularly important in densely populated deltas where ecological change and human vulnerability overlap.

7. Ecosystem-Specific Impacts

7.1 Mangrove Forests

Mangroves are highly productive intertidal forests that provide nursery habitat, fisheries support, shoreline stabilization and carbon storage. Their location makes them directly sensitive to sea-level rise, changes in sediment supply and storm regimes. They may initially respond to rising seas through vertical soil accretion and landward expansion, but this response becomes constrained when development prevents migration. UNEP emphasizes the ecological and livelihood importance of mangroves and the need for integrated conservation, restoration and sustainable management (UNEP, 2023).

7.2 Coral Reefs

Coral reefs are among the most climate-sensitive coastal ecosystems. Thermal stress causes bleaching, acidification reduces calcification potential, and storms can physically damage reef structures. Repeated disturbances reduce recovery time and can shift reefs from coral-dominated systems toward algae-dominated or structurally simplified states. The loss of reef complexity has implications for fish habitat, tourism and coastal protection. The IPCC concludes that almost all coral reefs face substantial degradation under future warming, with risks increasing sharply as warming rises (IPCC, 2022).

7.3 Seagrass Meadows and Salt Marshes

Seagrass meadows and salt marshes provide carbon storage, nursery habitat, sediment stabilization and water-quality functions. They are vulnerable to inundation, temperature stress, changing salinity, eutrophication and reduced water clarity. Salt marshes may migrate inland where space exists, but coastal infrastructure can prevent migration. Seagrass loss can also reduce sediment stability and carbon storage, creating feedbacks that amplify ecosystem degradation.

7.4 Estuaries, Beaches and Dunes

Estuaries are sensitive to changes in salinity, freshwater inflow, nutrient loading and temperature. Warming can increase the risk of hypoxia where nutrient enrichment is already high. Sandy beaches and dunes are affected by sea-level rise, erosion and storms, while development can limit their natural landward movement. The IPCC assesses increasing risks for estuaries and sandy beaches under higher levels of warming (IPCC, 2019).

8. International Comparative Case Studies

8.1 Great Barrier Reef, Australia

The Great Barrier Reef illustrates the cumulative effects of marine heatwaves, coral bleaching, storms, ocean acidification and local pressures. Repeated thermal disturbances can reduce coral cover and alter community composition. The case demonstrates that reef management can reduce local stressors and support recovery, but local conservation cannot fully compensate for continued global ocean warming. The international lesson is that local adaptation and global emissions reduction are complementary rather than interchangeable strategies.

8.2 Sundarbans, Bangladesh and India

The Sundarbans is the world's largest contiguous mangrove forest and a major natural buffer for a densely populated deltaic region. Climate-related sea-level rise, cyclones, salinity changes and erosion interact with sediment dynamics, upstream water management and human land use. The case highlights the importance of maintaining mangrove connectivity and allowing ecological

migration where possible. It also demonstrates that adaptation must include local livelihoods because ecosystem protection and social resilience are closely connected.

8.3 Mekong Delta, Viet Nam

The Mekong Delta demonstrates the compound nature of coastal vulnerability. Sea-level rise and salinity intrusion interact with land subsidence, sediment trapping, groundwater extraction, river engineering and changing agricultural practices. This illustrates an important principle of climate adaptation: reducing non-climatic pressures can increase the adaptive capacity of ecosystems and communities. Delta management therefore requires coordination across river basins, sectors and administrative boundaries.

8.4 Caribbean Coral-Reef Systems

Caribbean reefs have experienced bleaching, disease, storms, pollution, overfishing and habitat degradation. Climate change can intensify these pressures, reducing reef resilience. Regional experiences emphasize the value of herbivore protection, fisheries management, water-quality improvement, marine protected areas and reef restoration. However, restoration has ecological limits when thermal stress becomes too frequent or severe. This supports a two-level approach combining local reef stewardship with global climate mitigation.

8.5 Small Island Developing States

Small island developing states face a distinctive combination of sea-level rise, coastal flooding, coral-reef degradation, freshwater stress, extreme events and limited land availability. In many island settings, coral reefs, mangroves and seagrass meadows jointly contribute to coastal protection, fisheries and tourism. IPCC assessments indicate that even relatively small additional increments of sea-level rise can substantially increase flooding frequency in some island regions (IPCC, 2022). Adaptation therefore requires long-term planning, climate finance, ecosystem protection and, where necessary, carefully planned relocation.

9. Impacts on Ecosystem Services and Human Well-Being

Climate-driven degradation of coastal ecosystems affects society through multiple ecosystem-service pathways. Fisheries may decline when nursery habitats are damaged or species distributions shift. Tourism economies can be affected by coral bleaching, beach erosion and declining landscape quality. Mangrove and reef degradation can increase exposure to storm waves and coastal flooding. Coastal wetlands also store carbon, so their degradation can release previously stored carbon and reduce future sequestration potential.

The distribution of impacts is unequal. Communities with limited financial resources, high dependence on natural resources and restricted access to infrastructure or insurance may have fewer adaptation options. Indigenous peoples and small-scale fishing communities can also face cultural and livelihood losses that are not captured by conventional economic valuation. Effective adaptation therefore requires attention to distributional equity, participation and local knowledge in addition to physical risk reduction.

10. Adaptation and Nature-Based Solutions

10.1 Ecosystem-Based Adaptation

Ecosystem-based adaptation uses the protective and regulating functions of ecosystems to reduce climate risks. Mangrove restoration can attenuate waves and trap sediments; dunes can buffer storm impacts; and healthy coral reefs can reduce wave energy. Such approaches can provide biodiversity

and livelihood benefits in addition to hazard reduction. Their success depends on ecological suitability, sufficient space, long-term maintenance and protection from competing land uses.

10.2 Blue Carbon Conservation

Mangroves, salt marshes and seagrass meadows are commonly referred to as blue-carbon ecosystems because they can store substantial amounts of organic carbon in vegetation and sediments. The IPCC reports that restoration of vegetated coastal ecosystems could contribute to climate mitigation while delivering adaptation and biodiversity benefits (IPCC, 2019). However, blue-carbon initiatives should be based on robust carbon accounting and should not substitute for rapid reductions in fossil-fuel emissions.

10.3 Integrated Coastal Zone Management

Integrated coastal zone management brings ecological, economic and social considerations into a coordinated planning process. It can include setback zones, restoration areas, sustainable fisheries, pollution controls, flood-risk mapping, sediment management and climate-resilient infrastructure. Because coastal systems cross administrative boundaries, integration is needed among municipalities, national governments, river-basin authorities and neighboring countries where ecosystems or sediment and water flows are transboundary.

10.4 Planned Adaptation, Managed Retreat and Climate-Resilient Infrastructure

Hard infrastructure such as seawalls, dikes and surge barriers can reduce exposure in particular settings, but they can also alter sediment transport and disconnect wetlands from their natural migration pathways. Where protection becomes technically, economically or ecologically unsustainable, planned retreat may become part of the adaptation portfolio. The IPCC emphasizes that adaptation choices made today influence future options and that no single intervention can eliminate long-term sea-level-rise risk (IPCC, 2019).

11. International Governance and Policy Framework

Coastal climate resilience requires coordination among the global climate regime, biodiversity governance, disaster-risk reduction and sustainable development. The Paris Agreement provides the central framework for international climate mitigation and adaptation, while the Sustainable Development Goals connect climate action with poverty reduction, food security, sustainable cities and marine conservation. International biodiversity agreements and regional seas frameworks also contribute to ecosystem protection.

At the implementation level, international cooperation should prioritize accessible adaptation finance, scientific data sharing, technology transfer and capacity building. Small island and low-income coastal states often face high adaptation costs despite having limited historical responsibility for global greenhouse-gas emissions. A resilient international approach should therefore combine ambitious mitigation with predictable, equitable support for adaptation and loss-and-damage-related needs, consistent with applicable international agreements and decisions.

12. Discussion

The evidence reviewed in this study shows that climate change is transforming coastal ecosystems through interconnected physical, chemical and biological pathways. Three broad conclusions emerge. First, the pace of change matters as much as the absolute magnitude of change: ecosystems may adapt to gradual environmental shifts but struggle when hazards become rapid, recurrent or compound. Second, local management can strongly influence ecological resilience by reducing

pollution, overfishing, habitat fragmentation and destructive development. Third, there are limits to adaptation when global warming continues to intensify thermal stress and sea-level rise.

The international comparison also reveals that coastal resilience is context-dependent. A mangrove system with abundant sediment and undeveloped inland space may have more opportunity to migrate than a mangrove system enclosed by roads and settlements. A coral reef with low local pollution may recover more effectively after bleaching than a reef simultaneously affected by nutrient enrichment and overfishing. Therefore, international policy should not impose a single standardized adaptation model; it should provide principles, finance and scientific support that can be adapted to local ecological and social conditions.

An additional finding is that mitigation and adaptation should be treated as mutually reinforcing. Rapid emissions reductions reduce the rate of ocean warming, acidification and sea-level rise, thereby expanding the time available for adaptation. At the same time, ecosystem conservation can provide immediate resilience and carbon-storage benefits. The strongest policy portfolios therefore combine global mitigation, ecosystem protection, local risk reduction and inclusive governance.

13. Key Challenges and Research Gaps

- High uncertainty in local-scale projections of sea-level rise, sediment dynamics and extreme events.
- Insufficient long-term ecological monitoring in many low-income and small-island regions.
- Limited integration of ecological, socioeconomic and climate-risk datasets.
- Need for stronger evidence on the long-term effectiveness and scalability of restoration interventions.
- Difficulty valuing non-market ecosystem services, cultural values and distributional impacts.
- Governance fragmentation across coastal, river-basin, fisheries, urban-development and climate institutions.
- Need for improved climate finance access and locally led adaptation capacity.

14. Recommendations

- Accelerate greenhouse-gas mitigation because the long-term condition of coastal ecosystems is strongly constrained by the magnitude of global warming.
- Protect existing mangroves, salt marshes, seagrass meadows and coral reefs before investing primarily in restoration; intact ecosystems generally retain functions that are difficult to recreate.
- Restore degraded coastal ecosystems using site-specific ecological assessments, sediment and hydrological considerations, and long-term monitoring.
- Maintain or create landward migration corridors for coastal wetlands where sea-level rise threatens existing habitat.
- Integrate coastal climate risk into urban planning, infrastructure investment, fisheries management and disaster-risk reduction.
- Strengthen regional and transboundary cooperation for deltas, shared seas, migratory species and sediment systems.
- Expand equitable adaptation finance and technical support for vulnerable coastal states and communities.

- Combine scientific monitoring with Indigenous and local knowledge and meaningful community participation.
- Develop standardized indicators for ecosystem condition, resilience, adaptation effectiveness and ecosystem-service outcomes.
- Treat nature-based solutions as part of diversified adaptation portfolios rather than as universal substitutes for engineered protection or emissions reduction.

15. Conclusion

Global climate change is reshaping coastal ecosystems at a scale that has ecological, economic and social consequences across continents and ocean basins. Sea-level rise, ocean warming, marine heatwaves, acidification, changing storms and altered freshwater and sediment regimes are interacting with local pressures such as pollution, overfishing, habitat conversion and coastal development. The result is increasing stress on coral reefs, mangroves, seagrasses, salt marshes, estuaries, beaches and the communities that depend on them.

The international evidence demonstrates that coastal ecosystems are neither passive victims nor limitless buffers. They possess important adaptive capacities, but those capacities have ecological thresholds and are constrained by human development. Protecting space for wetland migration, reducing local stressors, restoring degraded habitats and maintaining ecological connectivity can strengthen resilience. Yet these measures cannot fully offset continued global warming. Consequently, effective coastal policy requires a dual strategy: rapid mitigation of greenhouse-gas emissions and sustained, locally appropriate adaptation.

The future of coastal ecosystems will depend not only on technological solutions but also on governance choices, finance, equity and international cooperation. A resilient coastal future requires policies that recognize ecosystems as infrastructure, communities as partners in adaptation, and climate mitigation as a prerequisite for maintaining ecological options. International research and policy should therefore move toward integrated, long-term and place-sensitive approaches capable of protecting both biodiversity and human well-being under changing climatic conditions.

References

- 1) Intergovernmental Panel on Climate Change (IPCC). (2018). Global warming of 1.5°C: An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels. <https://www.ipcc.ch/sr15/>
- 2) Intergovernmental Panel on Climate Change (IPCC). (2019). IPCC special report on the ocean and cryosphere in a changing climate. <https://www.ipcc.ch/srocc/>
- 3) Intergovernmental Panel on Climate Change (IPCC). (2022). Climate change 2022: Impacts, adaptation and vulnerability. Chapter 3: Oceans and coastal ecosystems and their services. Cambridge University Press. <https://www.ipcc.ch/report/ar6/wg2/chapter/chapter-3/>
- 4) Panel on Climate Change (IPCC). (2022). Climate change 2022: Impacts, adaptation and vulnerability. Chapter 15: Small islands. Cambridge University Press. <https://www.ipcc.ch/report/ar6/wg2/chapter/chapter-15/>
- 5) He, Q., Li, Z., Daleo, P., Lefcheck, J. S., Thomsen, M. S., Adams, J. B., & Bouma, T. J. (2025). Coastal wetland resilience through local, regional and global conservation. *Nature Reviews Biodiversity*, 1, 50–67. <https://www.nature.com/articles/s44358-024-00004-x>

- 6) Klein, S. G., Roch, C., & Duarte, C. M. (2024). Systematic review of the uncertainty of coral reef futures under climate change. *Nature Communications*, 15, 2224. <https://www.nature.com/articles/s41467-024-46255-2>
- 7) Nations. (n.d.). How is climate change impacting the world's ocean? United Nations Climate Change. <https://www.un.org/en/climatechange/science/climate-issues/ocean-impacts>
- 8) United Nations. (n.d.). Surging seas in a warming world. United Nations. <https://www.un.org/en/climatechange/reports/sea-level-rise>
- 9) United Nations Environment Programme (UNEP). (2023). Decades of mangrove forest change: What does it mean for nature, people and the climate? <https://www.unep.org/resources/report/decades-mangrove-forest-change-what-does-it-mean-nature-people-and-climate>
- 10) United Nations Environment Programme (UNEP). (2019). The impacts of climate change in coastal marine systems. <https://www.unep.org/index.php/resources/report/impacts-climate-change-coastal-marine-systems>
- 11) Zhang, Z., Luo, X., Friess, D. A., Wang, S., Li, Y., et al. (2024). Stronger increases but greater variability in global mangrove productivity compared to that of adjacent terrestrial forests. *Nature Ecology & Evolution*, 8, 239–250. <https://www.nature.com/articles/s41559-023-02264-w>

Appendix A. Comparative Summary of Major Coastal Climate Risks

Ecosystem	Major Climate Drivers	Principal Ecological Risks	Priority Responses
Mangroves	Sea-level rise; storms; salinity change; temperature	Inundation, coastal squeeze, habitat shifts	Protection, restoration, migration corridors, sediment management
Coral reefs	Marine heatwaves; warming; acidification; storms	Bleaching, mortality, reduced calcification, reef erosion	Emissions reduction, local stress reduction, targeted restoration
Seagrass	Warming; sea-level rise; turbidity; salinity change	Habitat loss, reduced productivity, carbon loss	Water-quality improvement, protection, restoration
Salt marshes	Sea-level rise; storms; salinity change	Inundation, vegetation shifts, coastal squeeze	Landward migration space, sediment management, restoration
Estuaries	Warming; altered rainfall; sea-level rise; nutrient loading	Hypoxia, salinization, biodiversity shifts	Integrated watershed management, pollution control
Beaches/dunes	Sea-level rise; storms; changing wave regimes	Erosion, habitat loss, infrastructure exposure	Setbacks, dune restoration, sediment management